Large Scale Structure: Can we deliver what we promised?

Shirley Ho, Carnegie Mellon University

LSSTC and LSST-DESC

Large Scale Structure Working Group Chair

Presenting results also from work within Planck and SDSS3-BOSS collaboration

Snowmass meeting, July 30th, 2013








Main Science goals


- Dark Energy: Constraining the expansion rate of the Universe via Baryon Acoustic Oscillations
- Ultimate test of non-gaussianities of the Universe? After Planck, large scale surveys is our best chance in probing fnl~I level non-gaussianities, we can do this with LSST!
- Total Neutrino masses: When combined with WMAP9, SDSS3 imaging data gives already a very stringent constraint of 0.26eV (95% CL)
- Dark Matter: We can probe modified gravity with redshift space distortions, and by combining with lensing, we can understand the bias of our tracers as a function of scale and redshift.

What are Baryon Acoustic Oscillations (BAO) anyway?

What are baryon acoustic oscillations (BAO)?

These fluctuations of 1 part in 10⁵ gravitationally grow into...





This sound wave can be used as a "standard ruler"


Dark energy changes this apparent ruler size

Courtesy slide from David Schlegel and animation from Daniel Eisenstein

What are Baryon Acoustic Oscillations (BAO) anyway?

What are baryon acoustic oscillations (BAO)?

These fluctuations of 1 part in 10⁵ gravitationally grow into...




...these ~unity fluctuations today

This sound wave can be used as a "standard ruler"


Dark energy changes this apparent ruler size

Courtesy slide from David Schlegel and animation from Daniel Eisenstein


Clustering in 3D




So we can do BAO in spectroscopic data like BOSS and SDSS!





AND we can do BAO in imaging data also (not only spectroscopic!)




AND we can do BAO in imaging data also (not only spectroscopic!)









AND we can do BAO in imaging data also (not only spectroscopic!)



So how do we do this Large Scale Structure measurement?





The First Challenge



The First Challenge


Make the measurement

Creating the sample, and the selection function properly

The First Challenge

Make the measurement


Creating the sample, and the selection function properly




SH, Cuesta, Seo, Ross, DePutter et al. (2012), Ross, SH et al. (2011)



Color offsets from Schlafly et al. 2011




The samples you can gleam from the sky is not a fair sample of the Universe, but rather a real Universe convolved with all the lovely observational systematics



Short summary:


BRUCE AND ASTRID MCWILLIAMS

Center for Cosmology



Ross, SH, Cuesta et al. (2011)


- Setting systematic weights for each galaxy, so that the effects of observational systematics are minimized (Ross, SH et al. 2011, Ross et al. 2012)
- Projecting out systematics by using systematic templates (Pullen & Hirata 2012, Slosar et al. 2013, Huterer et al. 2013)
- Cross-correlating observed densities with systematics to remove the effects of systematics on the over-densities (Myers et al. 2008, SH, Cuesta et al. 2012)
- Cross-correlating multiple redshift slices to detect unknown observational systematics (Agarwal, SH et al. 2013)




- Setting systematic weights for each galaxy, so that the effects of observational systematics are minimized (Ross, SH et al. 2011, Ross et al. 2012)
- Projecting out systematics by using systematic templates (Pullen & Hirata 2012, Slosar et al. 2013, Huterer et al. 2013)
- Cross-correlating observed densities with systematics to remove the effects of systematics on the over-densities (Myers et al. 2008, SH, Cuesta et al. 2012)
- Cross-correlating multiple redshift slices to detect unknown observational systematics (Agarwal, SH et al. 2013)

- Setting systematic weights for each galaxy, so that the effects of observational systematics are minimized (Ross, SH et al. 2011, Ross et al. 2012)
- Projecting out systematics by using systematic templates (Pullen & Hirata 2012, Slosar et al. 2013, Huterer et al. 2013)
- Cross-correlating observed densities with systematics to remove the effects of systematics on the over-densities (Myers et al. 2008, SH, Cuesta et al. 2012)
- Cross-correlating multiple redshift slices to detect unknown observational systematics (Agarwal, SH et al. 2013)

- Setting systematic weights for each galaxy, so that the effects of observational systematics are minimized (Ross, SH et al. 2011, Ross et al. 2012)
- Projecting out systematics by using systematic templates (Pullen & Hirata 2012, Slosar et al. 2013, Huterer et al. 2013)
- Cross-correlating observed densities with systematics to remove the effects of systematics on the over-densities (Myers et al. 2008, SH, Cuesta et al. 2012)
- Cross-correlating multiple redshift slices to detect unknown observational systematics (Agarwal, SH et al. 2013)



 Cross-correlating multiple redshift slices to detect unknown observational systematics (Agarwal, SH et al. 2013)



The Second Challenge:



e is

The Second Challenge: Beyond Linear Modeling

Yes, the theory of large scale structure is non-trivial:

The Second Challenge: Beyond Linear Modeling

Yes, the theory of large scale structure is non-trivial:

Observation:

redshift space galaxy correlation function

The Second Challenge: Beyond Linear Modeling


Yes, the theory of large scale structure is non-trivial:


Observation:

redshift space galaxy correlation function

What we can confidently predict:
real space dark matter correlation function


Current theory vs Simulations





Solid lines: Linear Theory

Points: N-body simulations



The Third Grand challenge

In orde
 matrix
 dumbes
 observat
 the vari

But this observa

Can we

riance easy and the whole ee what is s.

a large scale

The Third Grand challenge

- In order to generate reliable covariance matrix of our measurements, the easy and dumbest thing to do is to simulate the whole observation many many times, and see what is the variance across the simulations.
- But this can be Very expensive for a large scale observation such as LSST
- The second and third challenge can in principle be solved by increased ability to generate synthetic Universes


Generating Synthetic Universes

- The generation of synthetic Universe usually separated into two steps:
 - Creating Density field
 - Transformation of gaussian field (Coles 1992),
 - Perturbation theory like solution: PINNOCHIO (Monaco et al. 2002), 2LPT code (Scoccimaro & Sheth 2002), COLA (Tassev & Zaldarriaga 2013))
 - Painting galaxies onto the density field
 - Halo Occupation Distribution (Peacock & Smith 2002, Cooray & Sheth 2002, ..)
 - Semi-Analytical models (White 1978)


Other newer alternatives: Use Machine Learning to evolve the density field?




Very preliminary test: let's compare the halos statistics



Very preliminary: let's compare the halo statistics




Very preliminary: let's compare the halo statistics



If we can pass through all the challenges...

and there are more I didn't mention...


for example: photo-z determination, bias modeling, potential BAO fitting systematics ...



Robust Cosmological Constraints from LSS



Cosmological Constraints from the overall shape



Cosmological Constraints from the overall shape

